

Neck Dissection: Why, Which, When and How?

Dr Jeeve Kanagalingam
Consultant Head and Neck Surgeon,
Tan Tock Seng Hospital,
Singapore

***Choose well, cut well,
get well...***

History – radical neck

- Henry Butlin proposed enbloc removal of upper neck nodes with primary oral cavity cancers¹
- 'Radical' neck dissection first described by George Crile (1906)
- 60/132 patients enjoyed 3 year survival – 4 times better than control group²



¹ Butlin HI, Spencer WG, Disease of the tongue, 2nd ed. London: Cassell, 1900

² Crile G. Excision of cancer of the head and neck. With special reference to the plan of dissection based on 132 patients. JAMA 1906;47:1780–1786

History – selective neck

- Solis-Cohen proposed removal of uninvolved nodes during laryngectomy in 1901
- Functional Neck Dissection was described by Suarez in 1963¹
- Bocca popularised this, published outcome in 843 patients in 1984²

CLINICAL REVIEW

Changing patterns in the management of metastatic cancer to cervical lymph nodes have created a large number of surgical options. This historical review of the evolution of neck dissection offers a perspective that helps to interpret current standards and to anticipate future modifications.

Henry T. Hoffman, MD, *Section Editor*

FUNCTIONAL NECK DISSECTION: FACT AND FICTION

Alfio Ferlito, MD,¹ Javier Gavilán, MD,² J. Graham Buckley, MD,³
Ashok R. Shaha, MD, FACS,⁴ Adam J. Miodoński, MD, PhD,⁵ Alessandra Rinaldo, MD¹

¹ Department of Otolaryngology–Head and Neck Surgery, University of Udine, Policlinico Universitario, Piazzale S. Maria della Misericordia, 33100 Udine, Italy. E-mail: a.ferlito@dsc.uniud.it

² Department of Otorhinolaryngology, La Paz Hospital, Autonomous University, Madrid, Spain

³ Department of Otolaryngology–Head and Neck Surgery, Leeds General Infirmary, Leeds, UK

⁴ Head and Neck Service, Memorial Sloan-Kettering Cancer Center, New York, NY

⁵ Department of Otolaryngology, SEM Laboratory, Jagiellonian University, Kraków, Poland

Accepted 15 January 2001

Neck dissection is a valuable method for treating clinical, subclinical, and subpathologic metastasis from cancer of the head and neck. So much has been written and said on the subject that it is necessary to distinguish the fact from the fiction. If you think you have discovered something new, it is because you did not read enough. This popular statement is particularly valid for the neck dissection.

Jacob Da Silva Solis-Cohen (1838–1927) of Philadelphia, America's first head and neck surgeon,¹ mentioned the necessity of removing the lymphatics of the neck during total laryngectomy

Crile (1864–1943) of Cleveland, Ohio, developed "partial" and "radical" neck dissection operations, which involved removal of the primary tumor en bloc with the neck dissection.⁴ In 1906, he published the results of treatment of 132 head and neck cancers of which 60 patients enjoyed a 3-year survival—four times better than a comparable group with cervical metastases who did not have a neck dissection.³ This landmark article established the basis for effective treatment of such lesions by describing a block resection of the cervical lymph node-bearing tissue, either in continuity with the primary tumor or as a secondary operation for subsequent metastasis. The draw-

¹ Ferlito A et. al. Functional Neck Dissection: Fact and Fiction. *Head Neck* 2001;23:804-8

² Bocca E, Pignataro O, Oldini C, Cappa C. Functional neck dissection: an evaluation and review of 843 cases. *Laryngoscope* 1984;94:942–945

Why do a neck dissection?

- Eradicate disease

'When a single nodal metastasis exists at presentation or subsequently develops, the cure rate halves' ¹

- Stage the neck to guide further treatment and prognostic information
- Surgical access to primary tumour or for microvascular anastomosis

¹ Spiro RH, Alfonso AE, Farr HW, Strong EW. Cervical node metastases from epidermoid carcinoma of the oral cavity and oropharynx. A critical assessment of current staging. Am J Surg 1974;128:562-567.

Tumour biology

- Incidence of nodal metastases depends mainly on the site and the size of the primary tumour
 - 1% for early glottic tumours, 80% for nasopharyngeal carcinomas
- The majority of tumours will metastasise in a predictable manner but certain tumours will fast track to remote sites
 - nasopharyngeal cancers to level V
 - tongue cancers to jugulo-omohyoid nodes
 - pattern of spread will be disrupted by previous surgery or radiotherapy

Tumour biology

- Possibility of bilateral nodal disease should be considered especially when the primary site involves
 - Tongue base
 - Nasopharynx
 - Supraglottic larynx

Tumour biology

Oral cavity, anterior to circumvallate papillae



I-III, rarely to IV-V

Oropharynx



Mainly II, then III-IV, low rate to I, rarely V

Supraglottic larynx & hypopharynx



Mainly II, then III-IV, rarely I-V


Nasopharynx



Widespread II-V

Therapeutic dissections

n = 776




Shah JP, et al. The patterns of cervical lymph node metastases from squamous carcinoma of the oral cavity. *Cancer* 1990;66:109–13

Shah JP. Patterns of cervical lymph node metastasis from squamous carcinomas of the upper aerodigestive tract. *Am J Surg* 1990;160:405–9

Elective dissections


n = 343



Shah JP, et al. The patterns of cervical lymph node metastases from squamous carcinoma of the oral cavity. *Cancer* 1990;66:109–13

Shah JP. Patterns of cervical lymph node metastasis from squamous carcinomas of the upper aerodigestive tract. *Am J Surg* 1990;160:405–9

Oral cavity tumours



Spiro RH, Huvos AG, Wong GY, et al. Predictive value of tumor thickness in squamous carcinoma confined to the tongue and floor of the mouth. *Am J Surg* 1986;152: 345-50

Oral cavity: tumours > 4 mm thick predict mets

Original Article

Predictive Value of Tumor Thickness for Cervical Lymph-Node Involvement in Squamous Cell Carcinoma of the Oral Cavity

A Meta-analysis of Reported Studies


Shao Hui Huang, MSc^{1,2}, David Hwang, MB², Gina Lockwood, MMath³, David P. Goldstein, MD^{4,5},
and Brian O'Sullivan, MD^{2,4}

False negative rate of patients predicted by thickness not to have metastases rises significantly once tumour is more than 4 mm thick

Nasopharyngeal carcinoma

“an exception that proves the rule”

- 43 radical neck dissection specimens post-RT
 - In 70% there was more tumour bearing nodes than expected
 - 70% nodes involved had extra-capsular spread
 - 27.5% had tumour along XI nerve
 - 70% nodes were in posterior triangle



Wei WI et. al. Pathological basis of surgery in the management of postradiotherapy cervical metastasis in nasopharyngeal carcinoma. Arch Otolaryngol Head Neck Surg. 1992 Sep;118(9):923-9

Which neck dissection?

- Radical Neck Dissection
- Modified Radical Neck Dissection
 - Type I spare XI nerve
 - Type II spare XI and IJV
 - Type III spare XI, IJV and SCM
- Selective Neck Dissection
 - Supra-omohyoid (I-III)
 - Anterolateral (I-IV)
 - Lateral (II-IV)
 - Posterolateral (II-V)
 - Central (VI)

Robbins KT, Medina JE, Wolfe GT, Levine PA, Sessions RB, Pruet CW. Standardizing neck dissection terminology. Official report of the Academy's Committee for head and neck surgery and oncology. Arch Otolaryngol HeadNeck Surg 1991;117:601–605.

When to do a neck dissection?

Head and neck squamous cell carcinoma

■ N+ neck


- If primary disease is to be resected
- After definite radiotherapy, if there is residual nodal disease
- (After radiotherapy, if neck disease pre-treatment was bulky (N3) i.e. 'planned neck dissection')

■ N0 neck (elective neck dissection)

- If primary disease is to be resected and the rate of 'occult' metastases is 20% or more


Elective dissections

n = 343



Shah JP, et al. The patterns of cervical lymph node metastases from squamous carcinoma of the oral cavity. *Cancer* 1990;66:109–13

Shah JP. Patterns of cervical lymph node metastasis from squamous carcinomas of the upper aerodigestive tract. *Am J Surg* 1990;160:405–9



¹ Oral cavity - SOHND
 Oropharynx - LND or ALND
 Hypopharynx/larynx - LND

² Mandibulotomy, lateral pharyngotomy, supraglottic laryngectomy for example

³ Primary located in oropharynx, hypopharynx or supraglottic larynx. Oral cavity tumor greater than 2 mm thick

Why not observe the N0 neck closely?

60% of patients who recurred in the neck presented with N2 or greater disease

77% had evidence of extracapsular spread

Such patients required more extensive therapy than if they had undergone elective treatment

When to do a neck dissection?

Differentiated thyroid cancer

In N1a+, level VI (central compartment) neck dissection

In N1b+, level II-V (posterolateral) and level VI neck dissection

In N0 papillary thyroid cancer, if age > 45, male, >T2, offer elective level VI dissection



Medullary thyroid ca

In N0, level VI-VII neck dissection

In N0 and pT2-T4, or N1+ disease, add level IIa-Vb neck dissection

When to do a neck dissection

Salivary gland malignancy

If N+, modified radical neck dissection. XI may be difficult to preserve

If N0, consider level I-III and Va if high grade histology (e.g. high grade mucoepidermoid, undifferentiated, adenocarcinoma, SCC) T3-4, old age, SMG cancers and recurrent cancers



How to do a neck dissection


Operative technique

Preparation


- Position the patient appropriately with a shoulder roll
- Tape away hair, shave as necessary
- Give IV antibiotics no less than 30 minutes before first incision (Cefazolin)
- Plan your incision
- Infiltrate with 1 in 100,000 adrenaline




Apron with lazy 'S'




Hockey stick



Modified apron



McFee incision




Raising skin flaps

- Good retraction
- Hug the platysma and above veins
- Preserve Ext Jugular Vein for microvascular anastomosis
- In lateral neck dissection preserve Great Auricular Nerve
- Raise flaps to mandible, clavicle and trapezius for comprehensive necks





Raising subplatysmal flap



Identify and preserve marginal mandibular nerve

- Nerve is 1 cm inferior and posterior to the angle of the mandible
- Often retracted superiorly
- Deep to investing fascia but superficial to vein
- Hayes-Martin manoeuvre when oncologically safe





Lingual



Hypoglossal




For selective neck dissections

- 'Unsheath' the sternocleidomastoid
- Identify accessory nerve anteriorly
- Preserve C2 root to accessory
- Dissected level IIb

In all oral cavity and oropharyngeal tumours, in N+ larynx and hypopharyngeal tumours


- The posterior border of SCM and cervical plexus is the posterior limit of dissection


Unsheathing SCM





Accessory nerve


- Anteriorly nerve passes under branches of occipital artery to SCM
- Posteriorly Erb's point is a useful anatomical landmark
- Trace XI anteriorly under posterior belly of digastric













Stripping the IJV





Drains

- Place large suction drains
- Drain lateral gutters and submental and submandibular areas
- Watch for chylous drainage
- Remove once 'minimal drainage'



What can go wrong

- Skin flap necrosis
- Air embolus
- Bleeding
- Chyle leak
- Nerve damage


Post-op appearance



Thank you

Tumour biology

- Used oil-based medium injected pre-auricularly in 100 patients
- Nodal involvement examined in neck dissection specimens
- One-way only direction of lymphatic flow from level V to jugular chain



Tumour biology

- 2044 previously untreated HNSCC
- Location of metastases correlated with site and size of primary
- Clinical study and not pathological
- Findings later confirmed with pathological specimens by Byers
- 57% had cervical metastases
- Oral cavity, anterior tonsil pillar and soft palate, presence of metastases correlated with size of primary
- Tongue base, tonsil, supraglottic larynx and hypopharynx – no correlation

Lindberg R. Distribution of cervical lymph node metastases from squamous cell carcinoma of the upper respiratory and digestive tracts. *Cancer* 1972;29:1446–8

Byers RM, Wolf PF, Ballantyne AJ. Rationale for modified neck dissection. *Head Neck Surg* 1988;10:160–7.

